

SCHEDULING STATUS: Unscheduled

PROPRIETARY NAME (AND DOSAGE FORM):
EMEX SYRUP

COMPOSITION:

Each 5 ml contains:

Invert sugar	3,0 g
Phosphoric acid	25,0 mg
Methylparaben (preservative)	0,1% m/v

PHARMACOLOGICAL CLASSIFICATION:

A 5.7.2 Anti-emetics and anti-vertigo preparations

INDICATIONS:

Symptomatic relief of nausea in motion sickness and in early pregnancy.

CONTRA-INDICATIONS:

EMEX syrup must not be given to patients suffering from methyl alcohol poisoning, renal failure, diabetes and hereditary fructose intolerance.

WARNINGS:

EMEX syrup must not be taken for more than 1 hour (5 doses) without consulting a physician. If nausea continues or recurs frequently, consult a physician promptly since this may be a sign of a serious condition.

DOSAGE AND DIRECTIONS FOR USE:

Do not dilute the dosage since the optimal pH for functioning will be destroyed. Oral fluids should not be taken immediately before or for at least 15 minutes after the dose.

Motion sickness (car, air, etc.):

Adults: Three to six (15 – 30 ml) medicine measures. Dose may be repeated every fifteen minutes until distress subsides, but should not be taken for more than one hour (five doses) without consulting a physician.

Morning sickness: Three to six (15 – 30 ml) medicine measures. Dose should be taken on arising and repeated every three hours as needed.

Children over 3 years of age: One to two (5 – 10 ml) medicine measures. Dose may be repeated every

fifteen minutes until distress subsides, but should not be taken for more than one hour (five doses) without consulting a physician.

Children under 3 years of age: Use is not recommended.

SIDE-EFFECTS AND SPECIAL PRECAUTIONS:

EMEX produces no side-effects.

Special precautions: Fluids must not be taken immediately prior to or for at least 15 minutes after a dosage of EMEX because of the diluting effects.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

This is a safe and non-toxic medication.

IDENTIFICATION

A clear, colourless, spearmint-flavoured syrup.

PRESENTATION:

100 ml and 200 ml Amber glass bottle.

STORAGE INSTRUCTIONS:

Keep cool and protect from light. Store in a cupboard below 25 °C.

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER:

W/5.7.2/205

NAME AND BUSINESS ADDRESS OF APPLICANT:

Mirren (Pty) Ltd
18 Golden Drive
Morehill
Benoni
South Africa

DATE OF PUBLICATION OF THIS PACKAGE INSERT:

11 October 1999

SKEDULERINGSSTATUS: Nie geskeduleer nie

EIENDOMSNAAM (EN DOSEERVORM):

EMEX STROOP

SAMESTELLING:

Elke 5 ml bevat:

Invertsuiker	3,0 g
Fosforsuur	25,0 mg
Metielparabeen (Preserveermiddel)	0,1% m/v

FARMAKOLOGIESE KLASSIFIKASIE:

A 5.7.2 Anti-emetika en antivertigopreparate

FARMAKOLOGIESE INDIKASIES:

Simptomatiese verligting van naarheid by bewegingsiekte and vroeë swangerskap.

KONTRA-INDIKASIES:

EMEX stroop moet nie gegee word aan pasiënte wat ly aan metielalkoholvergiftiging, nierversaking, diabetes en erflike fruktose-intoleransie nie.

WAARSKUWING:

EMEX stroop moet nie vir meer as 1 uur (5 dosisse) geneem word sonder om 'n geneesheer te raadpleeg nie. As naarheid voortduur of dikwels terugkeer, raadpleeg 'n geneesheer onmiddellik aangesien dit 'n teken van 'n ernstige toestand kan wees.

DOSIS EN GEBRUIKSAANWYSINGS:

Moet nie verdun word nie want die optimale pH vir doeltreffende werking sal verander. Geen vloeistof moet onmiddellik voor of vir minstens 15 minute na die dosis gedrink word nie.

Bewegingsiekte (motor, lugreis, ens.):

Volwassenes: Drie tot ses (15 – 30 ml) medisynematevol. Dosis kan elke vyftien minute herhaal tot die naarheid verlig is, maar moet nie vir langer as 'n uur (5 doserings) gebruik word sonder om 'n geneesheer te raadpleeg nie.

Oggendsiekte: Drie tot ses (15 – 30 ml) medisynematevol. Dosering moet vroeg oggend (met opstaan) geneem word en daarna elke drie ure herhaal word, soos benodig.

Kinders ouer as 3 jaar: Een tot twee (5 – 10 ml) medisynematevol. Dosering kan elke vyftien minute herhaal word tot die naarheid verlig is, maar moet nie vir langer as 'n uur (5 doserings) gebruik word sonder om 'n geneesheer te raadpleeg nie.

Kinders jonger as 3 jaar: Gebruik word nie aanbeveel nie.

NEWE-EFFEKTE AND SPESIALE VOORSORGMATREËLS:

EMEX veroorsaak geen newe-effekte nie.

Spesiale voorsorgmaatreëls: Geen vloeistof moet onmiddellik voor of vir minstens 15 minute na 'n dosis EMEX gedrink word nie, omrede die verdunnende effek daarvan.

BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN:

EMEX is 'n veilige nie-toksiese medikasie.

IDENTIFIKASIE:

'n Helder, kleurlose stroop met 'n smaak van tuinment.

AANBIEDING:

100 ml en 200 ml Amber glas bottel.

BERGINGSAAWYSINGS:

Hou koel en beskerm teen lig. Bêre in 'n kas onder 25 °C.

HOU BUIE BEREIK VAN KINDERS.

REGISTRASIENOMMER:

W/5.7.2/205

NAAM EN BESIGHEIDSADRES VAN DIE APPLIKANT:

Mirren (Edms) Bpk
Goldenrylaan 18
Morehill
Benoni
Suid-Afrika

DATUM VAN PUBLIKASIE VAN HIERDIE VOUBILJET:

11 Oktober 1999