

SCHEDULING STATUS: Unscheduled

PROPRIETARY NAME (AND DOSAGE FORM):

ORANGO EFFERVESCENT TABLETS

COMPOSITION:

Each EFFERVESCENT TABLET contains:

| INGREDIENT | LABEL CLAIM | % RDA* |
|----------------------------------|--------------------|-----------------|
| BETACAROTENE (3 mg) | 500 µg RE | 50 |
| VITAMIN B1 | 15 mg | 1071 |
| VITAMIN B2 | 15 mg | 938 |
| VITAMIN B6 | 10 mg | 500 |
| VITAMIN B12 | 10 µg | 1000 |
| PANTOTHENIC ACID | 22,5 mg | 375 |
| VITAMIN C | 500 mg | 966 |
| NICOTINAMIDE | 24 mg | 133 |
| BIOTIN | 150 µg | 150 |
| FOLIC ACID | 200 µg | 100 |
| CALCIUM | 120 mg | 15 |
| MAGNESIUM | 100 mg | 33,3 |
| ZINC | 7,5 mg | 50 |
| SELENIUM (As Amino Acid Chelate) | 10,0 µg | Not Established |
| COPPER | 1,0 mg | Not Established |

*RDA = Recommended daily allowance for persons 4 years and older.

SUGAR FREE

Artificially sweetened with saccharin and cyclamate.

PHARMACOLOGICAL CLASSIFICATION:

A 22.1 Multivitamins and Multivitamins with Minerals.

PHARMACOLOGICAL ACTION:

Dietary Supplement.

INDICATIONS:

Orango Effervescent tablets are a supplement providing Vitamin C, the B group Vitamins, Betacarotene (Pro-vitamin A, Antioxidant) as well as the minerals Calcium, Magnesium, Selenium, Zinc and Copper.

CONTRA-INDICATIONS:

Renal impairment, dehydration and gastric lesions or disorders.

DOSAGE AND DIRECTIONS FOR USE:

Adults and children over 4 years of age:

Take one tablet, dissolved in a glass of water, daily.

SIDE-EFFECTS AND SPECIAL PRECAUTIONS:

Ascorbic acid (Vitamin C): Large doses may cause diarrhoea and the formation of renal calcium oxalate calculi. Daily doses of 600 mg or more have a diuretic action. Tolerance may be induced in patients taking high doses.

Diarrhoea, nausea and bowel discomfort may occur. Concurrent administration with zinc might diminish the effect of penicillamine. Tetracyclines may form a chelate with zinc.

Special precautions:

Pyridoxine hydrochloride (Vitamin B6) reduces the effects of levodopa. Cyanocobalamin (Vitamin B12) can mask symptoms of subacute degeneration of the spinal cord.

SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

The treatment of overdosage is symptomatic and supportive.

CONDITIONS OF REGISTRATION:

None.

IDENTIFICATION:

Large (25 mm), round, biplane, effervescent tablets. The tablets are light-yellow to grey with small dark-red particles impregnated into the tablet. The tablets effervesce in water with a slight foam. The red Betacarotene particles float to the surface at first, dissolving slowly to form an orange juice flavoured drink.

PRESENTATION:

Polypropylene tubes with 10 tablets, capped with a tamper-evident cap, containing drying material.

STORAGE INSTRUCTIONS:

Keep in a cool, dry place, below 25 °C.

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION/APPLICATION NUMBER:

To be allocated.

NAME AND ADDRESS OF THE APPLICANT:

Mirren (Pty) Ltd
18 Golden Drive
Morehill
Benoni
South Africa

DATE OF PUBLICATION:

10 August 2002

SKEDULERINGSSTATUS: Ongeskeduleerd

EIENDOMSNAAM (EN DOSEERVORM):
ORANGO BRUISTABLETTE

SAMESTELLING:

Elke BRUISTABLET bevat:

| BESTANDDEEL | PER TABLET | % ADT* |
|-------------------------------------|------------|------------------|
| BETA-KAROTEEN (3 mg) | 500 µg RE | 50 |
| VITAMIE B1 | 15 mg | 1071 |
| VITAMIE B2 | 15 mg | 938 |
| VITAMIE B6 | 10 mg | 500 |
| VITAMIE B12 | 10 µg | 1000 |
| PANTOTEENSUUR | 22,5 mg | 375 |
| VITAMIE C | 500 mg | 966 |
| NIKOTINAMIED | 24 mg | 133 |
| BIOTIEN | 150 µg | 150 |
| FOLIENSUUR | 200 µg | 100 |
| KALSIIUM | 120 mg | 15 |
| MAGNESIUM | 100 mg | 33,3 |
| SINK | 7,5 mg | 50 |
| SELEEN (As Aminosuur Chelaat) | 10,0 µg | Nie Vasgestel |
| KOPER | 1,0 mg | Nie Vasgestel |

*ADT = Aanbevole daaglikse dieettoelaag vir persone 4 jaar en ouer.

SUIKERVRY

Kunsmatig versoet met sakkarien en siklamaat.

FARMAKOLOGIESE KLASSIFIKASIE:

A 22.1 Multivitamie en Multivitamie met Minerale.

FARMAKOLOGIESE WERKING:

Dieetaanvulling.

INDIKASIES:

Orango Bruistablette is 'n suplement wat Vitamie C, Vitamie B kompleks, Beta-karoteen (Pro-vitamie A, Anti-oksidant) sowel as die minerale Kalsium, Magnesium, Seleen, Sink en Koper bevat.

KONTRA-INDIKASIES:

Renale ontoereikendheid, dehidrasie, gastriese letsels en wantoestande.

DOSIS EN GEBRUIKSAANWYSINGS:

Volwassenes en kinders ouer as 4 jaar:
Neem een tablet, opgelos in 'n glas water, daaglik.

NEWE-EFFEKTE EN SPESIALE

VOORSORGMATREËLS:

Askorbiensuur (Vitamie C): Groot dosisse mag diarree en die vorming van renale kalsium oksalaat kristalle by sekere individue veroorsaak. Dosisse van 600 mg en meer

per dag het 'n diuretiese effek. Toleransie mag geïnduseer word by individue wat groot dosisse oor 'n lang tydperk neem.

Diarree, naarheid en gastriese ongemak mag voorkom. Die werking van penisillamien word ingekort wanneer sink gelyktydig ingeneem word. Tetrasiklene vorm ook komplekse met sink.

Spesiale voorsorg:

Piridoksien (Vitamie B6) verminder die effek van Levodopa. Sianokobalamien (Vitamie B12) mag die simptome van sub-akute degenerasie van die spinale kolom maskeer.

SIMPOTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN:

Die behandeling van oordosering is simptomaties en ondersteunend.

VOORWAARDES VAN REGISTRASIE:

Geen.

IDENTIFIKASIE:

Orango is 'n groot (25 mm) ronde, bi-planêre bruistablet. Die tablet is liggeel tot grys met klein rooi spikkels in die tablet geïmpregneer. Die bruising in water is effens skuimend en die rooi betakaroteen deeltjies dryf eers bo-op die oplossing en los dan stadig op tot 'n aangename oranje-rooi lemoendrankie.

AANBIEDING:

Polipropileenbuis met 10 tablette. Die buise is geseël met 'n prop wat met 'n peuter-ring omsluit is. Binne-in die prop is drogingsmateriaal gesluit.

BERGINGSVOORSKRIFTE:

Bewaar in 'n koel, droë plek, onder 25 °C.
HOU BUITE BEREIK VAN KINDERS.

REGISTRASIE-NOMMER/

APPLIKASIE-NOMMER:

Moet nog toegeken word.

NAAM EN BESIGHEIDSADRES VAN

APPLIKANT:

Mirren (Edms) Bpk
Goldenrylaan 18
Morehill
Benoni
Suid-Afrika

DATUM VAN PUBLIKASIE:

10 Augustus 2002